

HIV

ACOI Board Review 2018
gerald.blackburn@beaumont.org

(No Disclosures)





David Kirby's mother, Kay, holds a photograph of her son -- taken by Ohio photographer Art Smith -- before AIDS took its toll.
Therese Frare




Bill Kirby tries to comfort his dying son, David, 1990.
Therese Frare

- 1.1 million HIV + in U.S.; (36.7 million worldwide)
 - 1 in 7 unaware of their HIV +; many others in denial
 - **responsible for up to 30% of transmission!**
 - < 1/3 completely virally suppressed
- 37,600 newly infected each year in U.S.; over 1/2 MSM
- 37,000 newly diagnosed each year will present with advanced disease (13,000 will die)

- Over half of HIV+ pts in U.S. are 50 y.o. or older; (by 2030, estimated ~73% will be over 50 y.o.) -> accelerated and/or increased incidence of:
 - CV Dx
 - Diabetes
 - Osteoporosis
 - COPD
 - Slower immune recovery
 - Malignancies
 - Other dx usually associated w/ aging, including cognitive disorders (or is it the meds?)
 - Drug interactions

HIV

- estimated to have entered the human population ~ 1920
- AIDS first described in U.S. in 1981; antibody testing first available 1985
- effective treatment first available in 1996; downside - tremendous pill burden, brutal side effects




Courtesy: AETC

- *irrevocable depletion of CD4 cells in GI tract and other lymphoid tissue
- without tx, approx. 10 yrs to develop AIDS
- initial presentation may be anywhere along this spectrum

The Acute Retroviral Syndrome

- Non-specific febrile illness often misdiagnosed as “**mono**” or “**aseptic meningitis**”, occurring 1 - 6 weeks following infection
 - chills, myalgias, adenopathy, maculopapular rash
 - pharyngitis, N/V, diarrhea
 - headache (LP-> mild pleocytosis) “aseptic meningitis”
 - elevated LFT's
- **Though HIV ab may be negative or indeterminate, these folks can be highly contagious** (if suspected, obtain HIV “Viral Load”)
- Spontaneous resolution over next few weeks

Window phase:
tests neg for HIV, but
highly contagious



Diagnosis

- Screening: EIA antibody (or other rapid tests)
- Testing now recommended as part of routine medical care (yearly if “high risk”). **CDC recs: yearly from ages 13 - 64**
- Newer assays that include p24 antigen (4th generation”) may be positive as early as 10 - 14 days after infection
- Confirmation: Western Blot
 - Time to positive: 4 - 5 weeks
 - Any two: p24, gp41, gp120/160 -> positive
 - one of above bands +, or other + bands -> “indeterminate”
 - if indeterminate, obtain quantitative assay for HIV by PCR - “viral load”

Clues to possible (untreated, advanced) HIV:

- Unusual presentation of a common illness
 - Pneumococcal pneumonia w/ bacteremia in a young person
 - Salmonella, shigella, campylobacter bacteremia
 - Severe or recurrent thrush, vaginal candidiasis
- Presentation of an unusual illness
 - Uncommon dx, e.g. cryptococcal meningitis
 - More advanced/severe dx than expected
 - Unusual age for illness
- TB, especially w/ unusual presentation
- Other STDs

Correlation of CD4 count to presentation of Opportunistic Infections/Malignancies

- Infections common in the non-HIV infected population tend to occur at higher CD4 counts.
As CD4 counts fall, these same infections may develop, but often with more extensive or disseminated disease. (TB, HSV-1 or 2, H. zoster, candidiasis)
- Infections rarely, if ever, seen in the non-immunosuppressed host tend to occur at the lowest CD4 counts e.g. disseminated CMV (100), MAC (50)
- Certain malignancies more common, even w/ “adequate” CD4 count

AIDS Defining Malignancies

- invasive cervical carcinoma
- Kaposi sarcoma
- systemic non-Hodgkin lymphoma
- primary CNS lymphoma




Non-AIDS Defining Malignancies Increased in HIV + Individuals

- lung
- liver
- kidney
- anus
- head & neck
- skin, including melanoma
- Hodgkin's lymphoma

O I's/neoplasms relative to CD4 counts

- **200 - 500 or above**
 - pulmonary TB
 - **bacterial pneumonia (pneumococcus most common)**
 - H. zoster
 - cervical CA, Kaposi's sarcoma, Hodgkin's lymphoma
 - oral/vaginal candidiasis; anemia; ITP; nephropathy (FSGS)



- < 200
 - PCP
 - Disseminated TB
 - Esophageal candidiasis
 - Cryptococcal meningitis; PML
 - Cryptosporidium
 - Non-Hodgkin's lymphoma
 - Disseminated histoplasmosis, coccidioidmycosis
 - wasting; dementia
- < 100
 - CNS toxo, lymphoma; disseminated CMV, MAC (elevated alk. phos)





CMV



Toxo: intensely white focal lesions w/ vitreous inflammation

Cryptococcal Meningitis

- Subacute, progressive headache; w/ or w/o fever
- Few to no WBC's in CSF
- + india ink, cryptococcal Ag
- Tx: Amphotericin B +/- flucytosine
by fluconazole
- T cell deficiencies




Still Common: PCP (PJP)

(*P. jirovecii*)

“YEE ROW VET ZEE”

- Subacute to acute pneumonia - still a common presentation in patients who are unaware of their HIV status or are otherwise untreated
- Diagnosis includes serum for: (1 -> 3) - beta - D - Glucan (Fungitell ®; Note: also used for invasive candidiasis and aspergillosis)®
- Tx: trimethoprim/sulfa; pentamidine if allergic
- **Steroids if $pO_2 < 70$**
(may not apply to HIV neg pts)



See JAMA: June 24, 2009

Warning:

- Multiple questions regarding trimethoprim/sulfa, including:
 - Side effects:
 - Maculopapular rash
 - Stevens-Johnson syndrome
 - TEN
 - Bone marrow suppression, other blood dyscrasias
 - Hyperkalemia
 - Volume overload w/ IV
 - Treatment of side effects
 - G6PD deficiency

Other Clues:
(if one STD, r/o others)





HSV



Kaposi's sarcoma (HSV-8)




Bacillary Angiomatosis (Bartonella sp.)

CMV Esophagitis

(D/D: CMV, HSV, Candida, aphthous ulcer)




Hairy Leukoplakia
(EBV)




GI Presentations

(Note: Tx will most always emphasize treating underlying HIV)

- Often chronic diarrheal syndrome
- Cryptosporidiosis - no fever; + AFB
 - AFB +
 - Cryptosporidium (3-6 microns)
 - also dx by direct immunofluorescence
 - Rx: ART; ? paromomycin, nitazoxanide
 - Cyclospora (7.5-10 microns)
 - Rx: TMP-SMX
 - Isospora (new: Cystoisospora)
 - Rx: TMP-SMX
 - Mycobacterium Avium-Complex "M.A.C."
 - Rx: Clarithro/Ethambutol/Rifabutin





- Other chronic diarrheal syndromes
 - Microsporidia (Enterocytozoon spp.) - no fever; bx w/ special stains
Rx: Tx HIV, albendazole
 - CMV - bloody diarrhea w/ fever; bx
Rx: ganciclovir
 - MAC - fever, wasting, diffuse abdominal pain; culture, +AFB
Rx: azithro or clarithro + ethambutol +/- rifabutin



Tx: Ganciclovir


Focal CNS syndromes

- Toxoplasmosis
 - Acute w/ multiple contrast + lesions w/ + serology (basal ganglia most often)
 - Fever
 - Mass effect





Focal CNS syndromes

- Lymphoma
(Usually diffuse, large B-cell)
- Subacute presentation
- Usually single contrast + lesion
- Mass effect
- Usually no fever
- + PCR for EBV (?)



Other Focal CNS syndromes

- PML
 - Multiple, contrast negative lesions of white matter
 - No mass effect
 - No fever
 - + PCR for JC virus



OI Prophylaxis*

PCP	CD4 < 200	TMP-SMX
TB	Previous + PPD** or +PPD > 5mm	INH x 9 mos (and others)
Toxoplasmosis	+ serology w/ CD4 < 100	TMP-SMX
M. avium complex	CD4 < 50	azithromycin or clarithromycin

*can usually be d/c'd upon return of CD4 count to above
threshold
parameters after ~3 months

**QFG assays similar to PPDs, but also w/ it's own difficulties in
interpretation

Clinical Course

- Viral load:
 - Correlates with degree of contagiousness, rate of immune deterioration (as reflected in CD4 ct)
 - “Cumulative viremia” w/ its resultant persistent inflammation and stimulation of the immune system may be responsible for many of the long term complications of HIV, e.g.,
 - Increased risk of CV and other diseases usually associated w/ aging
 - Increased risk of malignancy (including non-AIDS defining malignancies)

Clinical Course

- CD4 lymphocyte count (not the entire story):
 - Reflects immune status (as affected by VL)
 - Correlates w/ development of opportunistic infections (OI's)
 - Correlates to some extent w/ risk of malignancies, particularly if very low CD4 count prior to treatment
 - 27% of HIV-related deaths due to HIV-related malignancies
 - Risk of NHL > 76 times that of non-HIV infected individual
 - Restoration may approach normalcy, but probably never completely

Who/When to Treat?

(HHS Guidelines - Oct. 17, 2017)


Offer to all upon diagnosis, including
same day Rx

Benefits of Early Treatment

- Decrease in transmission (undetectable viremia minimizes risk of transmission)
- Decrease in illnesses associated w/ impaired immune system e.g. various infections, cancers
- Decrease in illnesses associated w/ chronic inflammation/accelerated aging e.g. heart disease, cancer

“Treatment as Prevention (of transmission)”

“No linked infections were observed when HIV-1 infection was stably suppressed by ART in the index participant”



Initiation of Antiretroviral Therapy in Early Asymptomatic HIV Infection

The INSIGHT START Study Group - N Engl J Med 2015; 373:797-807

- 4685 HIV + patients
 - median VL of 12,759 copies/ml
 - median CD4 count of 651
- After 3 years, those started immediately on ART experienced less than half of serious AIDS-related events (including reduced cancer risk by 64%) than those whose therapy was deferred to later

HIV and Pregnancy

- Overall risk of transmission if infected mother not identified (and not on tx): 25-33%
- IF infected mother identified (and appropriately treated): 1-3% or less
- Test at initial visit and at near term.
Treat if positive

Treatment

At least three drugs from at least two different classes of anti-retrovirals

- Usually **two NRTIs** (nucleoside reverse transcriptase inhibitor)

plus either a

- **NNRTI**

or a

- **PI/r** (ritonavir “boosted” protease inhibitor)

or a

- **InSTI** (integrase inhibitor)

Initial RX of Treatment-Naïve Patients HHS Guidelines, Oct. 2017

Integrase inhibitor based:

- Dolutegravir (Tivicay®) + abacavir/lamivudine = Triumeq® - ONLY for pts who are HLA-B*5701 negative or....
- Dolutegravir + [Emtricitabine/tenofovir (Truvada®)] or....
- Elvitegravir/cobistat + [Emtricitabine/tenofovir (Truvada®) / Descovy®] = Stribild® / Genvoya® or....
- Raltegravir (Isentress®) + [Emtricitabine/tenofovir (Truvada®)]

Note: Other sources have more extensive lists of acceptable options; certain two drug Rx's may be acceptable for long term suppression

Coinfection w/ Hepatitis B/C

- With hepatitis B:

Include combination of emtricitabine or lamivudine + tenofovir whenever possible, as these have dual activity for treating both infections

- Discontinuation may lead to serious liver damage from reactivation of Hepatitis B

- With hepatitis C:

most treat hepatitis C before initiating rx for HIV unless CD4 < 200

Possible ?'s:

- Hypersensitivity rxn to abacavir (Ziagen ®) if + for HLA-B*5701 (more common in caucasians).
DO NOT RX;
if prior reaction, DO NOT RE-CHALLENGE!!!
- Renal insufficiency, bone resorption w/ tenofovir (Viread ®) (or Truvada ® as combination Rx)
- Jaundice (indirect hyperbilirubinemia) w/ atazanivir (Reyataz ®)

Immune Reconstitution Inflammatory Syndrome (“IRIS”)

- An exaggerated inflammatory response to a previously relatively quiescent condition as a result of restoration of immune competence following initiation of HAART
 - Focal MAC
 - CMV vitreitis
 - TB
 - Cryptococcal meningitis
 - Hepatitis C
 - PML, HSV
- Rx: add anti-inflammatories and continue ART

Prevention:

- “Treatment as Prevention” - both of infection and complications of same
- Pre-exposure Prophylaxis (“PrEP”):
 - Once daily Truvada ®
 - Controversial, expensive, but effective if taken as rx'd
 - (Select) long term discordant sexual partners. Probably not necessary if partner undetectable VL
 - Commercial sex workers
 - but.....among MSM using PrEP:
 - 25.3 increased incidence of N. gonorrhoea!
 - 11.2 increased incidence of chlamydia!
 - 44.6 increased incidence of syphilis!!!

Prevention:

- Post-exposure Prophylaxis (“PEP”)
 - Occupational: Effective
 - Non-occupational (“nPEP”): at least partially effective
 - ~72 hr window for Rx
- Condoms; Circumcision

Potential ?'s

Acute Retroviral Syndrome

IRIS

Adverse Rxns to TMX/Sulfa

Presentations of HIV

Correlation of CD4 count w/ opportunistic infection

Histology of renal disease in HIV+ individuals: FSGS

Prophylaxis/Rx of OI's, e.g.:

Steroids in the treatment of PJP

Immune deficiency associated w/ Cryptococcal infections

TB prophylaxis / PPD skin test

References

- <http://www.aidsinfo.nih.gov> - Guidelines for the use of Antiretroviral Agents in HIV-1-Infected Adults and Adolescents. Oct. 17, 2017
- Cohen MS et al. Antiretroviral Therapy for the Prevention of HIV-1 Transmission. *N Engl J Med* 2016;375:830-839.
- Initiation of Antiretroviral Therapy in Early Asymptomatic HIV Infection. The INSIGHT START Study Group. *N Engl J Med* 2015; 373:797-807
- Grant et al. Preexposure chemoprophylaxis for HIV prevention in men who have sex with men. *NEJM* 2010; 363 (epub).
- Kitahata et al. Effect of Early versus Deferred Antiretroviral Therapy for HIV on Survival. *NEJM* 2009;360: 1815-1826
- Pilcher et al. Brief but Efficient: Acute HIV Infection and the Sexual Transmission of HIV. *JID* 2004;189: 1785-1792
- Zoufaly et al. Cumulative HIV Viremia during Highly Active Antiretroviral Therapy is a Strong Predictor of AIDS-Related Lymphoma. *JID* 2009;200: 79-87

Good Luck!