

# OFFICE MEDICINE FOR THE BOARDS

---

American College of Osteopathic Internists

Internal Medicine Board Review Course 2019

Baltimore Marriott Waterfront Hotel

Baltimore, Maryland

May 8- May 12, 2019

Scott Spradlin D.O. FACP,FACOI

I have no **relevant financial or nonfinancial relationships** in the products or services described, reviewed, evaluated or compared in this presentation.

# Objectives

- \* Review of Common Tests
- \* Review of Medical Ethics
- \* Review of Preventive Guidelines
- \* Review of Vaccinations
- \* Board Questions

# Common Tests

## CT Scans

Test of choice for:

- ✧ Cranial bleeds
- ✧ Renal lithiasis
- ✧ Appendicitis
- ✧ Diverticulitis

# Common Tests

## MR Scans

Test of choice for:

- ✧ CNS Diseases
- ✧ Avascular necrosis
- ✧ Aseptic necrosis
- ✧ Spinal Cord Diseases

# Common Tests

## V/Q Scan

- Positive = Segmental mismatched perfusion defect
- Pre-pneumonectomy- ok if FEV1 > 800cc

## Bone Scan

- Negative scan rules out osteomyelitis
- Very sensitive Paget Disease
- Best screening for Bone Metastasis
- Negative in Multiple Myeloma

## IVP

- To evaluate Obstructive concerns
- Remember CT to evaluate anatomic concerns

## Thyroid Scan

<sup>123</sup>I or <sup>131</sup>I

- Increased in Graves
- Decreased in Thyroiditis

## <sup>131</sup>I MIBG (iodine-131-meta-iodobenzylguanidine)

- Positive in Pheochromocytoma in 1-2 days

# Common Tests

Elevated Alkaline Phosphatase—  
”BLIP”

Bone

Late Pregnancy

Bone Metastasis

Pagets Disease

Osteomalacia

Liver

- Cholecystitis
- Primary Biliary Cirrhosis
- Primary Sclerosing Cholangitis
- Liver Disease

# Preoperative Evaluation

- For all surgeries, perioperative mortality is 0.3%
- Most perioperative deaths (55%) occur in the first 48 hours postoperatively
- Of perioperative deaths, 35% occur in the operating room, and 10% of deaths occur during anesthesia induction
- **Pulmonary complications are the most common perioperative complications**
- **Cardiac complications are the most common cause of perioperative death**
- Perioperative MIs usually occur by postoperative day 3
- Of perioperative MIs, 50% are fatal

# Surgical Clearance Review

## METs and Physical Activity

**METs, metabolic equivalents.**

### **1 MET**

Take care of self

Eat, dress, use toilet

Walk indoors around house

Walk 1–2 blocks on level ground at 2–3 mph

Dusting/washing dishes

(some classify this as 1–4 METs)

### **≥4 METs**

Climb 1 flight stairs or walk up a hill

Walk on level ground at 4 mph

Run a short distance

Scrubbing floors, moving heavy furniture

Golf, bowl, dance, doubles tennis, throw baseball or football

### **>10 METs**

Participate in strenuous sports including: singles tennis, football, basketball, skiing



# Surgical Clearance Review

## Surgery-Specific Risk

Low Risk (<1%)

Endoscopic procedures  
Superficial procedures  
Cataract surgery  
Breast surgery  
Ambulatory surgery

Intermediate Risk (<5%)

Carotid endarterectomy  
Endovascular abdominal  
aortic aneurysm repair  
Head and neck surgery  
Intraperitoneal surgery  
Intrathoracic surgery  
Orthopedic surgery  
Prostate surgery

High-Risk Vascular (>5%)

Peripheral vascular  
surgery  
Aortic/major vascular  
surgery

# Surgical Clearance Review

## **Active Cardiac Conditions (should Postpone or Cancel Surgery Until Resolved)**

Unstable coronary syndromes:  
Unstable angina  
Class III or IV angina  
Recent (i.e.,  $\leq 30$  days) myocardial infarction  
Decompensated heart failure  
Significant arrhythmia  
Supraventricular tachyarrhythmias with heart rate  $>100$   
High-grade atrioventricular block  
Mobitz II atrioventricular block  
Symptomatic ventricular arrhythmias  
Symptomatic bradycardia  
Severe valvular disease  
Severe aortic stenosis (i.e., mean pressure gradient  $>40$  mm Hg or aortic valve area  $<1$  cm, or symptomatic)  
Symptomatic mitral stenosis (i.e., progressive dyspnea on exertion, exertional syncope, or heart failure)

## **Clinical Risk Factors (should be Included in Preoperative Risk Assessment)**

History of heart disease  
History of compensated or prior heart failure  
History of cerebrovascular disease  
Diabetes mellitus (or specifically, anyone treated with insulin)  
Renal insufficiency (defined as serum creatinine  $>2$  mg/dL)

# Pre-surgery Appropriate Labs

- Hematocrit-
  - >65 years old
  - Blood loss surgeries
- Electrolytes
  - Not necessary unless known reason to monitor
- ECG
  - Vascular procedures
  - Major surgery
  - Cardiac disease/Hypertension/Diabetes
- CXR
  - Known respiratory disease or cardiac disease or >50
- PFT's
  - Unexplained respiratory issues.

# Summarizing Preoperative Risk

Preoperative risk is summarized by combining **clinical risk, functional status, and surgery-specific risk**

Patients with an **active cardiac condition should not undergo elective surgery** until the active cardiac condition has been treated

Patients with **no clinical risk factors can proceed to surgery without noninvasive cardiac testing**

Patients with **good functional status** can proceed to surgery **without noninvasive cardiac testing**

Patients undergoing **low-risk surgical procedures can proceed to surgery without noninvasive cardiac testing**

Patients with **one or two clinical risk factors undergoing intermediate-risk or vascular surgery or with poor or unknown functional status should be treated with perioperative  $\beta$ -blockers or consider noninvasive cardiac testing**

Patients with **three or more clinical risk factors should be managed similarly, unless undergoing vascular surgery, in which case they should undergo noninvasive cardiac testing**

# Medical Ethics

Beneficence:

*-Duty to act in the best interests of the patient*

Nonmaleficence:

*-Duty to do no harm*

Respect the patient's autonomy:

*-Duty to patient to make free choices*

Justice:

*-Duty to treat all patients fairly*

# Medical Ethics

## CPR/DNR:

- Whether or not CPR would be futile
- Preferences of the patient
- Expected quality of life

# Medical Ethics

## Brain Death:

- Physicians may stop tx if “brain dead”
- Ethics Committee review
- An EEG is not required for the diagnosis
- Organs can be donated without the patients prior consent

# Ethics

- Example:
- 55 year old man is diagnosed with cancer. As his treating physician you talk about the treatment medically but also discuss the social needs and support he will need. He asks you to not tell his wife?
- What are your obligations?
- What if it were a different diagnosis?



# Preventive Medicine

## Breast Cancer Screening

The U.S. Preventive Services Task Force (USPSTF) recommends screening **mammography**, with or without clinical breast examination (CBE), **every 1-2 years for women aged 50-74**.

The USPSTF concludes that the evidence is insufficient to recommend for or against routine CBE alone to screen for breast cancer.

The USPSTF concludes that the evidence is insufficient to recommend for or against teaching or performing routine breast self-examination (BSE).

# Preventive Medicine

## Screening for Cervical Cancer

The U.S. Preventive Services Task Force (USPSTF) strongly **recommends** screening for **cervical cancer in women ages 21-65 with cytology every 3 years or, women ages 30-65 who want to lengthen the screening with a combination of cytology and HPV testing every 5 years.**

USPSTF recommends against screening for cervical cancer in women <21.

The USPSTF recommends against cervical cancer screening in woman >65 who have had normal testing and not high risk

The USPSTF recommends **against** routine Pap smear screening in women who have had a total hysterectomy for benign disease.

The USPSTF concludes that the evidence is insufficient to recommend for or against the routine use of human papillomavirus (HPV) testing as a primary screening test for cervical cancer.

The USPSTF recommends against screening for cervical cancer with HPV alone or combination with cytology for women <30

# Preventive Medicine

## Colorectal Cancer

The U.S. Preventive Services Task Force (USPSTF) strongly recommends that clinicians screen men and women 50-75 years of age or older for colorectal cancer.

>75-85 Must have risk.

- FOBT/FIT DNA
- Barium enema/Sigmoidoscopy
- Colonoscopy
- **No evidence for colonography yet.**

# Preventive Medicine

## Screening for Ovarian Cancer

The U.S. Preventive Services Task Force (USPSTF) **recommends against routine screening for ovarian cancer.**

There is **NO** existing evidence that any screening test, including CA-125, ultrasound, or pelvic examination, reduces mortality from ovarian cancer.

# Preventive Medicine

## Screening for Abdominal Aortic Aneurysm

The U.S. Preventive Services Task Force (USPSTF) recommends one-time screening for abdominal aortic aneurysm (AAA) by ultrasonography in **men aged 65 to 75 who have ever smoked.**

The USPSTF recommends that clinicians *selectively screen* for screening for AAA in men aged 65 to 75 who have never smoked.

The USPSTF recommends **against** routine screening for AAA in women.

# Preventive Medicine

## Screening for Coronary Heart Disease

The U.S. Preventive Services Task Force (USPSTF) recommends against:

- ECG
- Exercise treadmill test (ETT)
- Electron-beam computerized tomography (EBCT)

For either the presence of severe CAS or the prediction of CHD events in adults at low risk for CHD events.

# Preventive Medicine

## High Blood Pressure

### Summary of Recommendations

The U.S. Preventive Services Task Force (USPSTF) strongly recommends that clinicians screen **adults aged 18 and older for high blood pressure.**

The USPSTF concludes that the evidence is insufficient to recommend for or against routine screening for high blood pressure in children and adolescents to reduce the risk of cardiovascular disease.

**At least every two years**

# Preventive Medicine

## Screening for Lipid Disorders in Adults

The U.S. Preventive Services Task Force (USPSTF) strongly recommends that clinicians routinely **screen men aged 35 years and older and women aged 45 years and older for lipid disorders and treat abnormal lipids in people who are at increased risk for coronary heart disease.**

The USPSTF recommends that clinicians routinely **screen younger adults (men aged 20 to 35 years and women aged 20 to 45 years) for lipid disorders if they have other risk factors for coronary heart disease.**

The USPSTF recommends that **screening for lipid disorders include measurement of total cholesterol (TC) and high-density lipoprotein cholesterol (HDL-C).**

USPSTF Statin Treatment recommends **low dose statin use** for:

Adults aged 40 to 75 years with no history of CVD, 1 or more CVD risk factors, and a calculated 10-year CVD event risk of 10% or greater. November 2016



# Preventive Medicine

## Screening for Chlamydial Infection

The U.S. Preventive Services Task Force (USPSTF) strongly recommends that clinicians routinely screen all sexually active **women aged 24 years and younger, and other asymptomatic women at increased risk for infection, for chlamydial infection.**

The USPSTF makes no recommendation for or against routinely screening asymptomatic low-risk women in the general population for chlamydial infection. The USPSTF **recommends that clinicians routinely screen all asymptomatic pregnant women aged 24 years and younger and others at increased risk for chlamydial infection.**

The USPSTF concludes that the evidence is insufficient to recommend for or against routinely screening asymptomatic men for chlamydial infection.

# Preventive Medicine

## Hormone Therapy for the Prevention of Chronic Conditions in Postmenopausal Women

The U.S. Preventive Services Task Force (USPSTF) recommends **against** the routine use of combined estrogen and progestin for the prevention of chronic conditions in postmenopausal women.

The U.S. Preventive Services Task Force (USPSTF) recommends **against** the routine use of unopposed estrogen for the prevention of chronic conditions in postmenopausal women who have had a hysterectomy.

# Preventive Medicine

## Screening for Type 2 Diabetes Mellitus in Adults

The U.S. Preventive Services Task Force (USPSTF) concludes that the **evidence is insufficient** to recommend for or against **routinely screening asymptomatic adults** for type 2 diabetes, impaired glucose tolerance, or impaired fasting glucose.

They do recommend screening for adults **40-70 years of age** who are **overweight or obese**

For certain **racial/ethnic groups** and family history should be screened

***FPG test (>126 mg/dL) / 2-hour PG level >200 mg/dL***

# Preventive Medicine

## Screening for Osteoporosis in Postmenopausal Women

The U.S. Preventive Services Task Force (USPSTF) **recommends that women aged 65 and older be screened routinely for osteoporosis.**

The USPSTF recommends for routine osteoporosis screening in postmenopausal women who are younger than 65 who are at increased risk of osteoporosis.

*For men the evidence is insufficient on the benefits and harms of screening for osteoporosis to prevent fracture without risks identified.*

# Preventive Medicine

## Aspirin for the Primary Prevention of Cardiovascular Events and Colorectal Cancer

The USPSTF recommends:

Low-dose aspirin use for the primary prevention of cardiovascular disease (CVD) and colorectal cancer (CRC) in adults aged **50 to 59 years** who have a 10% or greater 10-year CVD risk, are not at increased risk for bleeding, have a life expectancy of at least 10 years, and are willing to take low-dose aspirin daily for at least 10 years.

April 2016 addendum

# Preventive Medicine

## Screening for Lung Cancer

The USPSTF **recommends** annual screening for Lung Cancer with **low dose CT** in adults ages 55-80 who have a 30 pack year smoking history and currently smoke or have quit within 15 years.

Screening should be discontinued with negative scans and 15 year span since stopped smoking.

# Preventive Medicine

## Genetic Testing for BRCA

USPSTF recommends screening of woman who have family members with breast, ovarian, tubal, or peritoneal cancer with BRCA1 or BRCA2 susceptibility

If positive they should receive counseling and testing

# Preventive Medicine

## Carotid Artery Stenosis-screening

USPTF recommends **against** screening for asymptomatic carotid stenosis in the general population



# Preventive Medicine

## Hepatitis C

USPTF recommends screening for Hepatitis C (HCV) infection in persons at high risk for infection

USPTF also recommends offering 1-time screening for HCV infection to adults born 1945-1965

# Vaccines

## Live viruses

MMR, Polio(Oral), Nasal Flu, Yellow fever, Varicella, Rotovirus

## Live bacteria

BCG


## Dead viruses

Polio(IPV), Rabies, Flu, Hepatitis A, Pertussis, Hepatitis B

## Dead bacteria


Cholera, *H. influenza*, Pneumococcal, Meningococcal

# Images




# Images

- Macular Degeneration



# Images



# Images

- CMV Retinitis



# Images



# Images

- Toxoplasmosis






# Images




# Images

- Diabetic Retinopathy




# Images




# Images

- **Cholesterosis**




# Images




# Images

- Sickle Cell Retinopathy



# Images




# Images

- Metastatic Disease






# Images



# Images

- Retinal Vein Occlusion



# Ocular Eye Movement/CN

- Eye Muscles
- $LR_6(SO_4)_3$



# Dementia

- Progressive deterioration of cognition
- Insidious/Chronic
- No altered consciousness as in delirium
- 80 percent are presumed Alzheimers Disease
- To a lesser extent Multifarctional dementia/Lewy Body
- Common occurrence in Parkinsons
- Depression can resemble
  - Memory loss concern from family/friends
  - MMSE <24
  - Affect is normal
- Treatment
  - Cholinesterase inhibitors- Aricept/Exelon/Raadyne

# Dementia

- Lewy body-
  - Progressive cognitive decline
  - Hallucinations
  - Parkinsonian features
  - Sleep disturbances
- Pick's Disease-
  - Frontotemporal Dementia
  - Rapid cognitive decline
    - Language deficits
    - Apathy
    - Disinhibition
    - Emotional disturbances
  - CT/MRI – frontal and temporal atrophy
  - No definitive treatment

# Questions

- 19 yr old presents with a two day history of cough, runny nose, headache, and congestion. He has been having a low grade fever. Exam reveals T=99.4, nasal mucosa is red with yellowish secretions and the lungs are course but clear with cough.
- **Next best course of action would be?**
- A. Mucolytics, and acetaminophen
- B. Chest Radiograph, mucolytics, acetaminophen
- C. Macrolide antibiotic, mucolytics, and acetaminophen
- D. No action needed. It is a virus
- E. Amoxicillin, mucolytics, and acetaminophen

# Questions

- 19 yr old presents with a two day history of cough, runny nose, headache, and congestion. He has been having a low grade fever. Exam reveals T=99.4, nasal mucosa is red with yellowish secretions and the lungs are course but clear with cough.
- **Next best course of action would be?**
- **A. Mucolytics, and acetaminophen**
- B. Chest Radiograph, mucolytics, acetaminophen
- C. Macrolide antibiotic, mucolytics, and acetaminophen
- D. No action needed. It is a virus
- E. Amoxicillin, mucolytics, and acetaminophen

# Questions

- 60 yr old female comes in for her annual exam. PMH/PSH: Significant for HTN, Total Hysterectomy with BSO for benign disease in her 20's. She is on Benicar 40 for HTN. Allergies: NKDA.
- **Regarding a pap smear what is the next best course of action?**
- A. She is over 60 and has no need for a pap smear.
- B. She is post hysterectomy and doesn't need a pap smear.
- C. She needs a pap smear with HPV analysis.
- D. She needs a pap smear because we do not know the last three pap smears results with HPV analysis.
- E. She does need a pap smear because she is over 60 and I don't have the surgery report to confirm



# Questions

- 60 yr old female comes in for her annual exam. PMH/PSH: Significant for HTN, Total Hysterectomy with BSO for benign disease in her 20's. All her previous paps were negative. She is on Benicar 40 for HTN. Allergies: NKDA.
- **Regarding a pap smear what is the next best course of action?**
- A. She is over 60 and has no need for a pap smear.
- B. She is post hysterectomy and doesn't need a pap smear.
- C. She needs a pap smear with HPV analysis.
- D. She needs a pap smear because we do not know the last three pap smears results with HPV analysis.
- E. She needs a pap smear because she is over 60 and I dont have the surgical report

# Board Review Questions

- **Question 1**
- 32 yr old with history of intermittent back pain presents with **left leg pain**. The pain **starts in the lower back or buttocks area and radiates down the left leg into the foot**. He states that the pain seems to be worse at night. The patient is very healthy. He is an amateur bicycle racer and rides daily
- Normal PE except for the neuro exam. The spine is non-tender to palpation and percussion. The patient has moderate tenderness in the left buttocks area. It is difficult for the patient to tiptoe on his left foot. The rest of the neurological exam including deep tendon reflexes are normal.
- **Which of the following is the probable cause of this patient's pain?**
  - A. L5 radiculopathy
  - B. S1 radiculopathy
  - C. Sciatica
  - D. Peroneal nerve injury

# Board Review Questions

- **Question 1**

- 32 yr old with history of intermittent back pain presents with left leg pain. The pain starts in the lower back or buttocks area and radiates down the left leg into the foot. He states that the pain seems to be worse at night. The patient is very healthy. He is an amateur bicycle racer and rides daily
- Physical Exam
- Normal except for the neuro exam. The spine is non-tender to palpation and percussion. The patient has moderate tenderness in the left buttocks area. It is difficult for the patient to tiptoe on his left foot. The rest of the neurological exam including deep tendon reflexes are normal.
- Which of the following is the probable cause of this patient's pain?
  - A. L5 radiculopathy
  - B. S1 radiculopathy
  - **C. Sciatica**
  - D. Peroneal nerve injury

# Board Review Questions

- **Question 2**

- 34 year old moderately obese woman presents with complaints of having had an episode of **blindness in her left eye last week**. The symptoms **lasted approximately five hours** and then resolved spontaneously over the next couple hours. She has otherwise been in good health and does aerobic exercises three times a week. She notes that when she has **exercised since the episode she seems to have some dimming of vision in her left eye**. She is a nonsmoker and nondrinker. She has five children.
- Physical Exam
  - The pupil does not constrict when light is shone directly in her eye though consensual constriction is intact. She otherwise has a normal physical exam.
  - Lab: CBC, BUN, electrolytes, creatinine, and glucose are normal.
- Which of the following diagnoses is most consistent with this patient's signs and symptoms?
  - A. Optic Neuritis
  - B. Central retinal vein occlusion
  - C. Ischemic optic neuropathy
  - D. Pseudotumor cerebri
  - E. Acephalic migraine

# Board Review Questions

## Question 2

34 year old moderately obese woman presents with complaints of having had an episode of blindness in her left eye last week. The symptoms lasted approximately five hours and then resolved spontaneously over the next couple hours. She has otherwise been in good health and does aerobic exercises three times a week. She notes that when she has exercised since the episode she seems to have some dimming of vision in her left eye. She is a nonsmoker and nondrinker. She has five children.

### Physical Exam

The pupil does not constrict when light is shone directly in her eye though consensual constriction is intact. She otherwise has a normal physical exam.

Lab: CBC, BUN, electrolytes, Creatinine, and glucose are normal.

Which of the following diagnoses is most consistent with this patient's signs and symptoms?

- A. Optic Neuritis
- B. Central retinal vein occlusion
- C. Ischemic optic neuropathy
- D. Pseudotumor cerebri
- E. Acephalic migraine

# Board Review Questions

## Question 3

42 yr old man who is homeless and a chronic alcoholic is brought into the hospital by friends for evaluation of his impaired balance and decrease in mental status. On physical exam you find the patient has rather flat affect and has poor cognitive function. He has an ocular palsy involving the sixth cranial nerve.

What is the most probable cause of this patient's problem?

- A. Wernicke encephalopathy
- B. Korsakoff syndrome
- C. Subdural hematoma
- D. Subarachnoid bleed
- E. Thromboembolic stroke

# Board Review Questions

## Question 3

42 yr old man who is homeless and a chronic alcoholic is brought into the hospital by friends for evaluation of his impaired balance and decrease in mental status. On physical exam you find the patient has rather flat affect and has poor cognitive function. He has an ocular palsy involving the sixth cranial nerve.

What is the most probable cause of this patient's problem?

- A. Wernicke encephalopathy
- B. Korsakoff syndrome
- C. Subdural hematoma
- D. Subarachnoid bleed
- E. Thromboembolic stroke

# Board Review Questions

You are consulted on a case involving a previously healthy 34 yr old woman who presents with **complaints of weakness in her upper arms**. She says that the symptoms started two months previously and have slowly gotten worse. She works as a waitress and has noted that it has been especially difficult to put glasses in cabinets above her head. She complains of mild malaise.

On physical exam you find that the patient has mild tenderness of her upper arm and thigh area muscles. She has mildly decreased biceps, triceps, and deltoid strength. It is difficult to tell whether strength extinguishes with repeated testing because the patient is noncompliant.

Lab reveals CK 300 and ESR=60.

Electrolytes and CBC are normal.

What is the next most appropriate test?

- A. Edrophonium test
- B. Prednisone 10 mg per day
- C. Muscle Biopsy
- D. ANA,RA



# Board Review Questions

You are consulted on a case involving a previously healthy 34 yr old woman who presents with complaints of weakness in her upper arms. She says that the symptoms started two months previously and have slowly gotten worse. She works as a waitress and has noted that it has been especially difficult to put glasses in cabinets above her head. She complains of mild malaise. On physical exam you find that the patient has mild tenderness of her upper arm and thigh area muscles. She has mildly decreased biceps, triceps, and deltoid strength. It is difficult to tell whether strength extinguishes with repeated testing because the patient is noncompliant. Lab reveals CK 300 and ESR=60. Electrolytes and CBC are normal.

What is the next most appropriate test?

- A. Edrophonium test
- B. Prednisone 10 mg per day
- C. Muscle Biopsy
- D. ANA,RA

# Board Review Questions

**65 year old is brought in by her family to you for evaluation of abnormal gait, increased falling, and decreased mental status. Her family states that she seems to have extreme difficulty navigating the stairs. She also has been more forgetful than normal for the last three to six months. She is otherwise in good health.**

Physical Exam:

Elderly woman with a flat affect and masked facies.

She has tenseness in her neck and trunk.

Her neck is mildly hyperextended.

She has trouble looking down voluntarily.

No tremors are noted. Gait is slowed.

**Which of the following would be the most likely diagnosis for this patient?**

- A. Shy-Drager syndrome
- B. Parkinsons disease
- C. Progressive supranuclear palsy
- D. Autonomic dysfunction

**Which of the following provide good relief of symptoms in most of these patients?**

- A. Amantadine
- B. Deprenyl
- C. L-Dopa and carbidopa
- D. Bromocriptine
- E. No effective drug treatment therapy is available.

# Board Review Questions

65 year old is brought in by her family to you for evaluation of abnormal gait, increased falling, and decreased mental status. Her family states that she seems to have extreme difficulty navigating the stairs. She also has been more forgetful than normal for the last three to six months. She is otherwise in good health.

Physical Exam:

Elderly woman with a flat affect and masked facies.

She has tenseness in her neck and trunk.

Her neck is mildly hyperextended.

She has trouble looking down voluntarily.

No tremors are noted. Gait is slowed.

**Which of the following would be the most likely diagnosis for this patient?**

- A. Shy-Drager syndrome
- B. Parkinsons disease
- C. Progressive supranuclear palsy
- D. Autonomic dysfunction

**Which of the following provide good relief of symptoms in most of these patients?**

- A. Amantadine
- B. Deprenyl
- C. L-Dopa and carbidopa
- D. Bromocriptine
- E. No effective drug treatment therapy is available.

# Board Review Questions

62 year old presents with history of HTN and history of heart fluttering was brought into the ER with acute mental status change.

You have been called to evaluate by the ER doctor.

On exam she is not oriented to person or place or time.

BP 189/80. Heart is irregular and at a rate of 103.

Neurologically she is flaccid on the left upper extremity and the left lower extremity with a positive babinski.

She has a conjugate gaze to the right and aphasic.

**What is the diagnosis?**

**What is the distribution?**

**What area was affected?**



# Board Review Questions

62 year old presents with history of HTN and history of heart fluttering was brought into the ER with acute mental status change. You have been called to evaluate by the ER doctor.

On exam she is not oriented to person or place or time. BP 189/80. Heart is irregular and at a rate of 103. Neurologically she is flaccid on the left upper extremity and the left lower extremity with a positive babinski. She has a conjugate gaze to the right and aphasic

What is the diagnosis?

Right Ischemic Stroke

What is the distribution?

Middle Cerebral Artery-main trunk

What area was affected?

Middle cerebrum



# Board Question Review

50 year old with no medical history presents with weakness and fatigue.

No other history other than it has slowly got worse over the past few months.

On exam:

Noted fasciculations on exam of the chest and arms/legs and lower extremity weakness proximal greater than distal.

DTR's are +3/4 patellar and brisk.

Lab= CK 400

- What is the diagnosis?

# Board Question Review

50 year old with no medical history presents with weakness and fatigue. No other history other than it has slowly got worse over the past few months. On exam all is normal except noted fasciculations on exam of the chest and arms/legs and lower extremity weakness proximal greater than distal. DTR's are +3/4 patellar and brisk. Lab =CK 400

What is the diagnosis?

## Amyotrophic Lateral Sclerosis

- Most Common Anterior Horn Disease
- Lou Gehrig's Disease

# Image Challenge



**Q What is the diagnosis?**

**:**

- 1. Atlanto-occipital dislocation**
- 2. Atlanto-axial subluxation**
- 3. Pillar fracture**
- 4. Spinous process avulsion**
- 5. Wedge fracture**





# Image Challenge

**Q** What is the diagnosis?

:


**Answer:**

## **1. Atlanto-occipital dislocation**

This computed tomogram of the cervical spine shows major atlanto-occipital dislocation in the lateral view that proved to be fatal.



# Image Challenge



**Q What is the diagnosis?**

**:**

- 1. Central retinal vein occlusion**
- 2. Cholesterol embolism**
- 3. Retinoschisis**
- 4. Temporal arteritis**
- 5. Toxoplasmosis**



# Image Challenge

**Q What is the diagnosis?**

:

**Answer:**

## **2. Cholesterol embolism**

The image demonstrates cholesterol emboli with surrounding white lucency representing retinal edema. Fluorescein angiography confirmed occlusion of the cilioretinal artery with nonperfusion of the tissue bed in the hypofluorescent areas. Cholesterol emboli are crystals that are released in the arterial bloodstream from ulcerated or disrupted atherosclerotic plaques and can be an initial sign of vascular disease.



# Image Challenge



**Q** What is the diagnosis?

:

1. Lipoatrophy
2. Myositis ossificans
3. Rhabdomyolysis
4. Sarcoma
5. Syphilitic gumma



# Image Challenge

**Q What is the diagnosis?**

:


**Answer:**

## **1. Lipoatrophy**

Depressed areas of skin at the sites typically used for insulin injection are most consistent with injection-site lipoatrophy. It is associated with all types of insulin and may lead to erratic insulin absorption.



# Image Challenge



**Q What is the diagnosis?**

**:**

- 1. Brachial plexopathy**
- 2. Clavicular fracture**
- 3. Lipodystrophy**
- 4. Polyostotic fibrous dysplasia**
- 5. Rupture of the trapezius muscle**



# Image Challenge

**Q** What is the diagnosis?

:

**Answer:**

## **3. Lipodystrophy**

The presence of a dorsocervical fat pad in a patient who is otherwise lean suggests the diagnosis of lipodystrophy.



# Image Challenge



**Q** This patient presented with unilateral rhinorrhea. What is the diagnosis?

:

1. Nasal foreign body
2. Osteoma
3. Pituitary tumor
4. Skull fracture
5. Sinusitis





# Image Challenge

**Q** This patient presented with unilateral rhinorrhea. What is the diagnosis?

:


**Answer:**

## **3. Pituitary tumor**

The sinus radiograph reveals an enlarged sella turcica. Magnetic resonance imaging confirmed a pituitary macroadenoma that had caused cerebrospinal fluid rhinorrhea.



# Image Challenge



**Q** This technetium-99m sulfur colloid scan was performed after the patient presented with abdominal pain. Howell-Jolly bodies were present on a peripheral-blood smear. What is the diagnosis?

1. Hemochromatosis
2. Hydatidiform mole
3. Multiple myeloma
4. Pelvic spleen
5. Uterine sarcoma



# Image Challenge

**Q** This technetium-99m sulfur colloid scan was performed after the patient presented with abdominal pain. Howell-Jolly bodies were present on a peripheral-blood smear. What is the diagnosis?

**Answer:**

## **4. Pelvic spleen**

Abdominal imaging after injection of technetium-99m revealed a normal liver and a pelvic spleen. Pelvic spleen can be complicated by torsion of the pedicle. Howell-Jolly bodies suggest functional asplenia.



# Image Challenge



**Q** This patient with a history of rheumatoid arthritis presented with a several-month history of a painful left calf. What is the diagnosis?

- 1. Baker's cyst**
- 2. Gastrocnemius tear**
- 3. Meniscal cyst**
- 4. Lipoblastoma**
- 5. Septic arthritis**



# Image Challenge

**Q** This patient with a history of rheumatoid arthritis presented with a several-month history of a painful left calf. What is the diagnosis?


**Answer:**

## 1. Baker's cyst

Baker's cysts may be seen in patients with rheumatoid arthritis, in whom they consist of a synovium-lined sac that is continuous with the joint space. Symptoms of a Baker's cyst, especially if it ruptures, may mimic those of venous thrombosis. Meniscal cysts appear laterally. Lipoblastomas typically affect young children. There are no signs to suggest septic arthritis. The presentation is not typical of a gastrocnemius tear.



# Image Challenge



**Q What is the diagnosis?**

**:**

- 1. Cytomegalovirus retinitis**
- 2. Glaucoma**
- 3. Malignant hypertension**
- 4. Panretinal photocoagulation**
- 5. Retinal vein thrombosis**



# Image Challenge

**Q What is the diagnosis?**

:

**Answer:**

## **3. Malignant hypertension**

The retinal image demonstrates disk swelling with scattered cotton-wool spots, and inferior serous retinal detachment. The patient's blood pressure was 220/150 mm Hg. The pattern of subretinal fluid accumulation and exudative retinal detachment is most consistent with malignant hypertension.



# Image Challenge



**Q What physical findings would be expected to be present in this patient?**

**:**

- 1. Quadriplegia with bilateral gaze paresis**
- 2. Left hemiparesis, gaze deviated to the left**
- 3. Left hemiparesis, gaze deviated to the right**
- 4. Right hemiparesis, gaze deviated to the left**
- 5. Right hemiparesis, gaze deviated to the right**





# Image Challenge

**Q What physical findings would be expected to be present in this patient?**

:

**Answer:**

**3. Left hemiparesis, gaze deviated to the right**

The images demonstrate an infarct of the right internal capsule nucleus which caused left hemiparesis and gaze deviation to the right.



# Image Challenge



**Q What is the diagnosis?**

**:**

- 1. Hemorrhagic episcleritis**
- 2. Intravitreal hemorrhage**
- 3. Loa loa**
- 4. Osteogenesis imperfecta**
- 5. Traumatic bleb**



# Image Challenge

**Q What is the diagnosis?**

:

**Answer:**

## **5. Traumatic bleb**

Contact ultrasonographic biomicroscopy of this large pigmented lesion in the left eye revealed a cystic structure with communication to the vitreous cavity, prompting the diagnosis of a uveal bleb rather than melanoma.



## Image Challenge



**Q Which one of the following signs would you expect to find in this 76-year-old woman with anemia?**

- 1. Bulbar palsy**
- 2. Flaccid paralysis of the upper extremities**
- 3. Lower-extremity spasticity**
- 4. Lower-extremity thermoanesthesia**
- 5. Romberg's sign**



# Image Challenge

**Q** Which one of the following signs would you expect to find in this 76-year-old woman with anemia?


**Answer:**

## **5. Romberg's sign**

The MRI reveals hyperintense signal in the dorsal spinal cord extending from C2 to C5. Vibratory and proprioception are carried in the dorsal column, and were decreased in this woman's fingers and toes. Romberg's sign was present. She was unable to perform tandem gait. Pernicious anemia was diagnosed.



# Image Challenge



**Q** What term is used to describe this finding?

:

1. Arc eye
2. Asthenopia
3. Choroideremia
4. Coloboma
5. Corectopia



# Image Challenge

**Q** What term is used to describe this finding?

:


**Answer:**

## **4. Coloboma**

Colobomas are the result of abnormal closure of the optic fissure. They may occur anywhere along the optic fissure and can affect the iris, choroid, or macula. Isolated iris colobomas are asymptomatic, but those involving the macula or the optic disk can result in severe visual impairment. Typical iris colobomas occur in the inferonasal quadrant.



# Image Challenge



**Q What clinical presentation would be expected in this patient?**

**:**

- 1. Asymmetrical mydriasis**
- 2. Ataxic hemiparesis**
- 3. Hypothermia**
- 4. Quadriplegia**
- 5. Upward gaze palsy**





# Image Challenge

**Q** What clinical presentation would be expected in this patient?

:


**Answer:**

## **4. Quadriplegia**

The most common presentation of a pontine hemorrhage is quadriplegia. Small, reactive pupils are characteristic of pontine hemorrhages. Hemiparesis would be expected if the hemorrhage were asymmetrical. Hypothermia is unusual. Upward gaze palsy occurs with midbrain involvement.



# Image Challenge



**Q What is the diagnosis?**

**:**

- 1. Epidural hematoma**
- 2. Glioblastoma multiforme**
- 3. Meningioma**
- 4. Subarachnoid hemorrhage**
- 5. Subdural hematoma**



# Image Challenge

**Q** What is the diagnosis?

:

**Answer:**

## 1. Epidural hematoma

Computed tomogram shows a 2.5-cm epidural hematoma in the left parietal region with mass effect, effacement, and left-to-right midline shift. Epidural hematomas have a lens-shaped appearance. Subdural hematomas are typically sickle-shaped.



# Image Challenge



**Q This patient with chronic alcoholism presented with dysarthria and horizontal nystagmus. What is the diagnosis?**

- 1. Brainstem glioma**
- 2. Central pontine myelinolysis**
- 3. Neurosarcoidosis**
- 4. Pontine stroke**
- 5. Tabes dorsalis**



# Image Challenge

**Q** This patient with chronic alcoholism presented with dysarthria and horizontal  
: nystagmus. What is the diagnosis?

**Answer:**

## **2. Central pontine myelinolysis**

The brain MRI reveals central pontine myelinolysis with a well-defined lesion in the pons of low T1-signal intensity, with sparing of the ventral lateral and cortical spinal tracts and no space-occupying effect or distortion of the adjacent fourth ventricle. Central pontine myelinolysis is a noninflammatory, demyelinating condition that was originally described in those with chronic alcoholism.



# Image Challenge



**Q** What diagnosis is suggested by this corneal photograph?

:

1. Anterior uveitis
2. Chlamydia trachomatis infection
3. Cytomegalovirus retinitis
4. Herpes simplex virus infection
5. Toxocariasis



# Image Challenge

**Q** What diagnosis is suggested by this corneal photograph?

:


**Answer:**

**4. Herpes simplex virus infection**

A dendritic ulcer is indicative of herpes simplex virus infection.



# Image Challenge



**Q Which one of the following patterns of visual disturbance would be predicted to be demonstrable on examination of this patient?**

- 1. Inferior hemifield loss**
- 2. Temporal quadrantanopsia**
- 3. Unilateral blindness**
- 4. Macular sparing hemianopia**
- 5. Peripheral ring scotoma**





# Image Challenge

**Q Which one of the following patterns of visual disturbance would be predicted to be demonstrable on examination of this patient?**

**Answer:**


## **1. Inferior hemifield loss**

Dilated ophthalmoscopy of this left eye illustrates a nonrefractile plaque in the proximal superior retinal artery with retinal whitening in the superior macula signifying retinal ischemia. Superior retinal ischemia will result in a defect in the inferior visual field, as in this patient. A diagnosis of hemiretinal arterial occlusion was made.

*Read More: N Engl J Med 2008;358:2716*



# Image Challenge



**Q** What term is used to describe this finding?

:

1. Hyphema
2. Hypopyon
3. Iridocyclitis
4. Iridodonesis
5. Synechia



# Image Challenge

**Q** What term is used to describe this finding?

:

**Answer:**

## 1. Hyphema

Layering of blood in the anterior segment is termed hyphema. Hypopyon refers to pus in the anterior segment. Iridocyclitis refers to inflammation of the iris. Iridodonesis is a quivering of the iris when the patient moves the eye. Synechia is an adhesion between the iris and the lens.

*Read More: N Engl J Med 2008;358:2265*



# Image Challenge



**Q** What is the diagnosis?

:

1. Central retinal artery occlusion
2. Diabetic papillopathy
3. Ocular toxoplasmosis
4. Optic neuritis
5. Malignant hypertension



# Image Challenge

**Q What is the diagnosis?**

:

**Answer:**

## **5. Malignant hypertension**

The fundus photograph shows disk edema, cottonwool spots, a swollen optic nerve, and retinal hemorrhages. Together, these findings suggest a diagnosis of malignant hypertension.

*Read More: N Engl J Med 2008;358:1951*



# Image Challenge



**Q** A patient with this tomogram would be most likely to present with which one of the following signs?

- 1. Unilateral blindness**
- 2. Hemiplegia**
- 3. Alexia without agraphia**
- 4. Hemiballismus**
- 5. Internuclear ophthalmoplegia**



# Image Challenge

**Q** A patient with this tomogram would be most likely to present with which one of the following signs?

**Answer:**

## **2. Hemiplegia**

The tomogram shows a calcified object in the proximal right middle cerebral artery. Occlusion of the middle cerebral artery would be most likely to be associated with contralateral hemiparesis, as in this case. The other listed choices represent stroke syndromes that most typically involve other vascular territories.



# Image Challenge



**Q What is the most likely diagnosis?**

**:**

- 1. Paget's disease**
- 2. Meningioma**
- 3. Neurocysticercosis**
- 4. Pneumocephalus**
- 5. Hyperparathyroidism**





# Image Challenge

**Q** What is the most likely diagnosis?

:

**Answer:**

## **4. Pneumocephalus**

The cranial radiograph shows air in the left temporal region without evidence of fracture. Pneumocephalus can occur after neurosurgical procedures, head and facial trauma, or ear infection and can even occur spontaneously. The finding is not typical of a brown tumor, Paget's disease, neurocysticercosis, or meningioma.

*Read More: N Engl J Med 2008;358:e13*



# Image Challenge



**Q This patient is most likely to have presented with which one of the following findings?**

**:**

- 1. Amnesia**
- 2. Ataxia**
- 3. Dysphagia**
- 4. Hemianopia**
- 5. Hemiparesis**



# Image Challenge

**Q** This patient is most likely to have presented with which one of the following findings?  
:

**Answer:**


## **5. Hemiparesis**

An acute occlusion of the middle cerebral artery, as illustrated here, is most likely to present with contralateral hemiparesis.

*Read More: N Engl J Med 2007;357:2495*



# Image Challenge



**Q** What is the diagnosis?

:

1. Hypertriglyceridemia
2. Hypertensive retinopathy
3. Optic atrophy
4. Central retinal artery occlusion
5. Cytomegalovirus retinitis



# Image Challenge

**Q What is the diagnosis?**

:

**Answer:**

## **1. Hypertriglyceridemia**

The creamy white vessels in the fundus resulted from extreme hypertriglyceridemia.

*Read More: N Engl J Med 1999;340:1969*



# Image Challenge



**Q What is the diagnosis?**

**:**

- 1. Intracranial hemorrhage**
- 2. Osteoma**
- 3. Neurocysticercosis**
- 4. Arachnoid cyst**
- 5. Meningioma**



# Image Challenge

**Q** What is the diagnosis?

:

**Answer:**

## **5. Meningioma**

This well-circumscribed and highly calcified extra-axial mass is most consistent with a meningioma.

*Read More: N Engl J Med 2007;356:e14*



# Image Challenge



**Q What is the diagnosis?**

**:**

- 1. Chalazion**
- 2. Papilloma**
- 3. Pterygium**
- 4. Pinguecula**
- 5. Coloboma**





# Image Challenge

**Q What is the diagnosis?**

:

**Answer:**

## **2. Papilloma**

This 9-year-old boy was diagnosed with conjunctival viral papilloma.

*Read More: N Engl J Med 2007;356:1352*



# Image Challenge



**Q This patient developed difficulty swallowing following a dental procedure. What is the diagnosis?**

- 1. Spondylolisthesis**
- 2. Prevertebral air**
- 3. Pharyngeal diverticulum**
- 4. Pharyngeal foreign body**
- 5. Periodontal abscess**



# Image Challenge

**Q** This patient developed difficulty swallowing following a dental procedure. What is the diagnosis?

**Answer:**

## **2. Prevertebral air**

The radiograph shows emphysema with prevertebral air in the cervical soft tissues. A high-speed dental drill was implicated.



# Image Challenge



**Q** These lesions were neither pruritic nor painful. What is the diagnosis?

:

1. Pyoderma gangrenosus
2. Phlegmasia cerulea dolens
3. Pretibial myxedema
4. Necrobiosis lipoidica diabetorum
5. Erythema nodosum



# Image Challenge

**Q** These lesions were neither pruritic nor painful. What is the diagnosis?

:

**Answer:**

## **4. Necrobiosis lipoidica diabetorum**

This patient was diagnosed with necrobiosis lipoidica diabetorum.

*Read More: N Engl J Med 2006;355:e20*



# Image Challenge



**Q What is the diagnosis?**

**:**

- 1. Cytomegalovirus retinitis**
- 2. Roth spots**
- 3. Central retinal vein occlusion**
- 4. Hypertensive retinopathy**
- 5. Papilledema**



# Image Challenge

**Q** What is the diagnosis?

:

**Answer:**


## **5. Papilledema**

The fundoscopic image suggests florid papilledema.

*Read More: N Engl J Med 2006;355:1262*



## Image Challenge



**Q** A 55-year-old kidney-transplant recipient presented with headache and fever. The cerebrospinal fluid contained 84 percent neutrophils. What is the most likely diagnosis?

1. *Nocardia asteroides* infection
2. Cerebral toxoplasmosis
3. *Listeria monocytogenes* infection
4. Miliary tuberculosis
5. *Cryptococcus neoformans* infection





# Image Challenge

**Q** A 55-year-old kidney-transplant recipient presented with headache and fever. The  
**:** cerebrospinal fluid contained 84 percent neutrophils. What is the most likely diagnosis?

**Answer:**

## **1. Nocardia asteroides infection**

*Nocardia asteroides* infection typically presents with an abscess or multiple enhancing lesions, with neutrophilic pleocytosis in the cerebrospinal fluid.

*Read More: N Engl J Med 2006;354:2802*



# Image Challenge



**Q What is the diagnosis?**

**:**

- 1. Left facial palsy**
- 2. Cavernous sinus thrombosis**
- 3. Orbital lymphoma**
- 4. Herpes zoster ophthalmicus**
- 5. Orbital fracture**



# Image Challenge

**Q** What is the diagnosis?

:

**Answer:**

## **4. Herpes zoster ophthalmicus**

The image illustrates a left VI nerve palsy from herpes zoster ophthalmicus.

*Read More: N Engl J Med 2005;353:e14*



# Good Luck!!

- ✓ Relax
- ✓ Take A Deep Breath
- ✓ Read the Question
- ✓ Think on what was presented- *nothing else*
- ✓ Make the Decision
- ✓ Pass the exam for certification/recertification
- ✓ Congratulations!
- ✓ Thank you for your commitment to Medicine